

IIT-JEE 2008 EXAMINATION PAPER

(SOLUTIONS)

PAPER – I

Part – I (MATHEMATICS)

SECTION – I

Straight Objective Type

13/04/08

This section contains 6 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

1. The total number of local maxima and local minima of the function


$$f(x) = \begin{cases} (2+x)^3, & -3 < x \leq -1 \\ x^{2/3}, & -1 < x < 2 \end{cases} \text{ is -}$$

- (A) 0
(C) 2

- (B) 1
(D) 3

[Ans. C]

Sol.



2. Let $g(x) = \frac{(x-1)^n}{\log \cos^m(x-1)}$; $0 < x < 2$, m and n are integers, $m \neq 0$, $n > 0$, and let p be the left hand derivative of $|x-1|$ at $x=1$. If $\lim_{x \rightarrow 1^+} g(x) = p$, then

- (A) $n=1, m=1$
(C) $n=2, m=2$

- (B) $n=1, m=-1$
(D) $n > 2, m=n$

[Ans. C]

Sol. $p =$ left hand derivative of $|x-1|$
 $= -1$

$$\lim_{x \rightarrow 1^+} g(x) = p$$

$$= -1$$

$$\lim_{x \rightarrow 1^+} \frac{(x-1)^n}{\log \cos^m(x-1)} = -1$$

$$\lim_{x \rightarrow 1^+} \frac{n(x-1)^{n-1}}{\frac{-m \cos^{m-1}(x-1) \cdot \sin(x-1)}{\cos^m(x-1)}} = -1$$

$$\lim_{x \rightarrow 1^+} \frac{n(x-1)^{n-1}}{-m \tan(x-1)} = -1$$

if $n-1 = 1 \Rightarrow n = 2$

and $m = 2.$

3. Let a and b be non-zero real numbers. Then, the equation $(ax^2 + by^2 + c)(x^2 - 5xy + 6y^2) = 0$ represents
- (A) four straight lines, when $c = 0$ and a, b are of the same sign
 (B) two straight lines and a circle, when $a = b$, and c is of sign opposite to that of a
 (C) two straight lines and a hyperbola, when a and b are of the same sign and c is of sign opposite to that of a
 (D) a circle and an ellipse, when a and b are of the same sign and c is of sign opposite to that of a

[Ans. B]

Sol. On solving, $x^2 - 5xy + 6y^2 = 0$ gives two lines passing through origin and $a = b, c = -a$ satisfies to the curve $ax^2 + by^2 + c = 0$ for circle.

4. The edges of a parallelepiped are of unit length and are parallel to non-coplanar unit vectors $\hat{a}, \hat{b}, \hat{c}$ such that $\hat{a} \cdot \hat{b} = \hat{b} \cdot \hat{c} = \hat{c} \cdot \hat{a} = 1/2$. Then, the volume of the parallelepiped is

(A) $\frac{1}{\sqrt{2}}$ (B) $\frac{1}{2\sqrt{2}}$ (C) $\frac{\sqrt{3}}{2}$ (D) $\frac{1}{\sqrt{3}}$

[Ans. A]

Sol. Volume of a parallelepiped is $[abc]$

$$\therefore \bar{a} \cdot \bar{b} = |a| |b| \cos \theta = \frac{1}{2} \Rightarrow \theta = 60^\circ$$

Now,

$$\text{Volume} = |[\hat{a} \hat{b} \hat{c}]|$$

$$= |(\hat{a} \times \hat{b}) \cdot \hat{c}| = \sqrt{\begin{vmatrix} \hat{a} \cdot \hat{a} & \hat{a} \cdot \hat{b} & \hat{a} \cdot \hat{c} \\ \hat{b} \cdot \hat{a} & \hat{b} \cdot \hat{b} & \hat{b} \cdot \hat{c} \\ \hat{c} \cdot \hat{a} & \hat{c} \cdot \hat{b} & \hat{c} \cdot \hat{c} \end{vmatrix}} = \frac{1}{\sqrt{2}}$$

5. If $0 < x < 1$, then $\sqrt{1+x^2} [\{x \cos(\cot^{-1} x) + \sin(\cot^{-1} x)\}^2 - 1]^{1/2} =$

(A) $\frac{x}{\sqrt{1+x^2}}$

(B) x


(C) $x\sqrt{1+x^2}$

(D) $\sqrt{1+x^2}$

[Ans. C]

Sol. $\sqrt{1+x^2} [\{x \cos(\cot^{-1} x) + \sin(\cot^{-1} x)\}^2 - 1]^{1/2}$

$$= \sqrt{1+x^2} \left[\left\{ x \cdot \frac{x}{\sqrt{1+x^2}} + \frac{1}{\sqrt{1+x^2}} \right\}^2 - 1 \right]^{1/2}$$



$$= \sqrt{1+x^2} \left[\left(\frac{x^2}{\sqrt{1+x^2}} + \frac{1}{\sqrt{1+x^2}} \right)^2 - 1 \right]^{1/2}$$

$$= \sqrt{1+x^2} [x^2 + 1 - 1]^{1/2}$$

$$= x\sqrt{1+x^2}$$

6. Consider the two curves

$$C_1 : y^2 = 4x$$

$$C_2 : x^2 + y^2 - 6x + 1 = 0$$

Then,

(A) C_1 and C_2 touch each other only at one point

(B) C_1 and C_2 touch each other exactly at two points

(C) C_1 and C_2 intersect (but do not touch) at exactly two points

(D) C_1 and C_2 neither intersect nor touch each other

[Ans. B]

Sol. On solving the given two curves C_1 and C_2 we get the points of tangency $(1, \pm 2)$.

SECTION – II

Multiple Correct Answers Type

This section contains 4 multiple correct answer(s) type questions. Each question has 4 choices (A), (B), (C) and (D) out of which ONE OR MORE is/are correct.

7. A straight line through the vertex P of a triangle PQR intersects the side QR at the point S and the circumcircle of the triangle PQR at the point T. If S is not the centre of the circumcircle, then

(A) $\frac{1}{PS} + \frac{1}{ST} < \frac{2}{\sqrt{QS \times SR}}$ (B) $\frac{1}{PS} + \frac{1}{ST} > \frac{2}{\sqrt{QS \times SR}}$
 (C) $\frac{1}{PS} + \frac{1}{ST} < \frac{4}{QR}$ (D) $\frac{1}{PS} + \frac{1}{ST} > \frac{4}{QR}$

[Ans. B, D]

Sol. By using A.M \geq G.M. inequality, we get the answers.

8. Let

$$S_n = \sum_{k=1}^n \frac{n}{n^2 + kn + k^2} \text{ and } T_n = \sum_{k=0}^{n-1} \frac{n}{n^2 + kn + k^2}$$

for $n = 1, 2, 3, \dots$. Then,

(A) $S_n < \frac{\pi}{3\sqrt{3}}$ (B) $S_n > \frac{\pi}{3\sqrt{3}}$ (C) $T_n < \frac{\pi}{3\sqrt{3}}$ (D) $T_n > \frac{\pi}{3\sqrt{3}}$

[Ans. A,C]

Sol. $T_n = \sum_{k=0}^{n-1} \frac{n}{n^2 + kn + k^2}$

$$\lim_{n \rightarrow \infty} T_n = \int_0^1 \frac{dx}{1+x+x^2} \qquad \text{Let } x = \frac{k}{n}$$

$$= \int_0^1 \frac{dx}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}}$$

$$\begin{aligned}
&= \frac{2}{\sqrt{3}} \tan^{-1} \frac{x + \frac{1}{2}}{\frac{\sqrt{3}}{2}} \Big|_0^1 \\
&= \frac{2}{\sqrt{3}} \tan^{-1} \frac{2x+1}{\sqrt{3}} \Big|_0^1 = \frac{2}{\sqrt{3}} \left[\tan^{-1} \sqrt{3} - \tan^{-1} \frac{1}{\sqrt{3}} \right] \\
&= \frac{2}{\sqrt{3}} \left[\frac{\pi}{3} - \frac{\pi}{6} \right]
\end{aligned}$$

$$\lim_{n \rightarrow \infty} T_n = \frac{2}{\sqrt{3}} \cdot \frac{\pi}{6} = \frac{\pi}{3\sqrt{3}} \quad \text{This is for } n \rightarrow \infty$$

\therefore But $n < \infty$

$$\therefore T_n < \frac{\pi}{3\sqrt{3}}$$

$$S_n = \sum_{k=1}^n \frac{n}{n^2 + kn + k^2}$$

$$\lim_{n \rightarrow \infty} S_n = \int_0^1 \frac{dx}{1+x+x^2} \quad \text{Let } x = \frac{k}{n}$$

$$= \frac{\pi}{3\sqrt{3}} \quad \text{Again } n \rightarrow \infty \text{ then } S_n = \frac{\pi}{3\sqrt{3}}$$

$$\therefore S_n < \frac{\pi}{3\sqrt{3}} \text{ as } n \text{ is not exactly } \infty.$$

9. Let $f(x)$ be a non-constant twice differentiable function defined on $(-\infty, \infty)$ such that $f(x) = f(1-x)$ and

$$f'\left(\frac{1}{4}\right) = 0. \text{ Then,}$$

(A) $f''(x)$ vanishes at least twice on $[0, 1]$

(B) $f'\left(\frac{1}{2}\right) = 0$

(C) $\int_{-1/2}^{1/2} f\left(x + \frac{1}{2}\right) \sin x \, dx = 0$

(D) $\int_0^{1/2} f(t) e^{\sin \pi t} \, dt = \int_{1/2}^1 f(1-t) e^{\sin \pi t} \, dt$

[Ans. A,B,C,D]

Sol. $f(x) = f(1-x)$

$$f'(x) = -f'(1-x) \Rightarrow f'(1/4) = -f'(3/4) = 0 \text{ (given)}$$

put $x = 1/2$

$$\Rightarrow f'\left(\frac{1}{2}\right) = -f'\left(\frac{1}{2}\right)$$

$$\Rightarrow f'\left(\frac{1}{2}\right) = 0$$

Since $f'(1/2) = f'(1/4) = f'(3/4) = 0$ hence by Rolle's theorem statement A is true.

as given $f(x) = f(1-x)$

replace x by $x + \frac{1}{2}$

$$f\left(\frac{1}{2} + x\right) = f\left[1 - \left(\frac{1}{2} + x\right)\right] = f\left(\frac{1}{2} - x\right) \quad \dots(1)$$

$$\text{Now, } I = \int_{-1/2}^{1/2} f\left(x + \frac{1}{2}\right) \sin x \, dx$$

$$I = - \int_{-1/2}^{1/2} f\left(-x + \frac{1}{2}\right) \sin x \, dx \quad (\text{put } x = -x)$$

$$I = - \int_{-1/2}^{1/2} f\left(\frac{1}{2} + x\right) \sin x \, dx \quad [\text{by (1)}]$$

$$I = -I$$

So $I = 0$

Now, let $t = 1 - x$

$$- \int_1^{1/2} f(1-x) e^{\sin \pi x} \, dx$$

$$= \int_{1/2}^1 f(1-t) e^{\sin \pi t} \, dt$$

10. Let $P(x_1, y_1)$ and $Q(x_2, y_2)$, $y_1 < 0$, $y_2 < 0$, be the end points of the latus rectum of the ellipse $x^2 + 4y^2 = 4$.

The equations of parabolas with latus rectum PQ are

(A) $x^2 + 2\sqrt{3}y = 3 + \sqrt{3}$

(B) $x^2 - 2\sqrt{3}y = 3 + \sqrt{3}$

(C) $x^2 + 2\sqrt{3}y = 3 - \sqrt{3}$

(D) $x^2 - 2\sqrt{3}y = 3 - \sqrt{3}$

[Ans. B,C]

Sol. Let ellipse is $\frac{x^2}{4} + \frac{y^2}{1} = 1$.

P and Q points are $\left(\pm\sqrt{3}, -\frac{1}{2}\right)$, focus S of the ellipse is $\left(0, -\frac{1}{2}\right)$. Now find directrix of the parabola

parallel to PQ and at $\sqrt{3}$ unit distance from PQ, and then use $PS^2 = PM^2$ to find the parabola.

SECTION – III
Reasoning Type

This section contains 4 reasoning type questions. Each question has 4 choices (A), (B), (C) and (D), out of which ONLY ONE is correct.

11. Consider three planes

$$P_1 : x - y + z = 1$$

$$P_2 : x + y - z = -1$$

$$P_3 : x - 3y + 3z = 2.$$

Let L_1, L_2, L_3 be the lines of intersection of the planes P_2 and P_3 , P_3 and P_1 , and P_1 and P_2 , respectively.

STATEMENT - 1

At least two of the lines L_1, L_2 and L_3 are non-parallel

and

STATEMENT - 2

The three planes do not have a common point

(A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1

(B) Statement-1 is True, Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1

(C) Statement-1 is True, Statement-2 is False

(D) Statement-1 is False, Statement-2 is True.

[Ans. D]

Sol. The given three planes intersect on parallel lines.

12. Let f and g be real valued functions defined on interval $(-1, 1)$ such that $g''(x)$ is continuous, $g(0) \neq 0$, $g'(0) = 0$, $g''(0) \neq 0$, and $f(x) = g(x) \sin x$.

STATEMENT - 1

$$\lim_{x \rightarrow 0} [g(x) \cot x - g(0) \operatorname{cosec} x] = f''(0)$$

and

STATEMENT - 2

$$f'(0) = g(0).$$

(A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1

(B) Statement-1 is True, Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1

(C) Statement-1 is True, Statement-2 is False

(D) Statement-1 is False, Statement-2 is True.

[Ans. B]

Sol. Statement 1. $\lim_{x \rightarrow 0} \frac{g(x)\cos x - g(0)}{\sin x}$

$$= \lim_{x \rightarrow 0} \frac{g'(x)\cos x - g(x)\sin x}{\cos x} \quad (\text{Applying L - H rule})$$

$$= g'(0) - 0 = 0 = f'(0) \quad (\text{True})$$

Statement 2

$$f'(x) = g(x)\cos x + g'(x)\sin x$$

$$f'(0) = g(0) \quad (\text{Ture})$$

13. Consider the system of equations

$$\begin{aligned} x - 2y + 3z &= -1 \\ -x + y - 2z &= k \\ x - 3y + 4z &= 1 \end{aligned}$$

STATEMENT -1 : The system of equations has no solution for $k \neq 3$.

and

STATEMENT -2 : The determinant $\begin{vmatrix} 1 & 3 & -1 \\ -1 & -2 & k \\ 1 & 4 & 1 \end{vmatrix} \neq 0$, for $k \neq 3$.

(A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1

(B) Statement-1 is True, Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1

(C) Statement-1 is True, Statement-2 is False

(D) Statement-1 is False, Statement-2 is True.

[Ans. A]

Sol. since $\Delta = \begin{vmatrix} 1 & -2 & 3 \\ -1 & 1 & -2 \\ 1 & -3 & 4 \end{vmatrix} = 0$

\therefore for having either $\Delta_x \neq 0$ or $\Delta_y \neq 0$ or $\Delta_z \neq 0$ no solution

$$\therefore \Delta_x = \begin{vmatrix} -1 & -2 & 3 \\ k & 1 & -2 \\ 1 & -3 & 4 \end{vmatrix} \neq 0$$

$$\Rightarrow 3 - k \neq 0$$

$$\Rightarrow k \neq 3$$

Now again

$$\begin{vmatrix} 1 & 3 & -1 \\ -1 & -2 & k \\ 1 & 4 & 1 \end{vmatrix} \neq 0$$

$$\Rightarrow k \neq 3$$

14. Consider the system of equations

$$ax + by = 0, cx + dy = 0, \text{ where } a, b, c, d \in \{0, 1\}$$

STATEMENT -1 : The probability that the system of equations has a unique solution is $\frac{3}{8}$.

and

STATEMENT-2 : The probability that the system of equations has a solution is 1.

(A) Statement-1 is True, Statement-2 is True; Statement-2 **is** a correct explanation for Statement-1

(B) Statement-1 is True, Statement-2 is True; Statement-2 **is NOT** a correct explanation for Statement-1

(C) Statement-1 is True, Statement-2 is False

(D) Statement-1 is False, Statement-2 is True.

[Ans. B]

Sol. Total possibilities = $2^4 = 16$

Now favourable solution is possible if $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = 0$

\therefore cases possible are

$$\begin{vmatrix} 1 & 1 \\ 0 & 0 \end{vmatrix}, \begin{vmatrix} 0 & 0 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 1 & 0 \\ 1 & 0 \end{vmatrix}, \begin{vmatrix} 0 & 1 \\ 0 & 1 \end{vmatrix}, \begin{vmatrix} 0 & 0 \\ 0 & 0 \end{vmatrix}, \begin{vmatrix} 1 & 1 \\ 1 & 1 \end{vmatrix} = 6$$

$$\text{So probability} = \frac{6}{16} = \frac{3}{8}$$

SECTION – IV

Linked Comprehension Type

This section contains 3 paragraphs. Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D), out of which ONLY ONE is correct.

Paragraph for Question Nos. 15 to 17

Let A, B, C be three sets of complex numbers as defined below

$$A = \{z : \text{Im}z \geq 1\}$$

$$B = \{z : |z - 2 - i| = 3\}$$

$$C = \{z : \text{Re}((1 - i)z) = \sqrt{2}\}$$

15. The number of elements in the set $A \cap B \cap C$ is

(A) 0

(B) 1

(C) 2

(D) ∞


[Ans.B]

Sol. Given that,

$$A : y \geq 1$$

$$B : (x - 2)^2 + (y - 1)^2 = 9$$

$$C : x + y = \sqrt{2}$$



There is only one common point which satisfies to all the three given curves

16. Let z be any point in $A \cap B \cap C$. Then, $|z + 1 - i|^2 + |z - 5 - i|^2$ lies between

(A) 25 and 29

(B) 30 and 34

(C) 35 and 39

(D) 40 and 44

[Ans.C]

Sol. Let $z_1 = (-1, 1)$ and $z_2 = (5, 1)$.

$$|z + 1 - i|^2 + |z - 5 - i|^2 = |z_1 - z_2|^2 = (6)^2 = 36.$$

17. Let z be any point in $A \cap B \cap C$ and let w be any point satisfying $|w - 2 - i| < 3$.

Then, $|z| - |w| + 3$ lies between

(A) -6 and 3

(B) -3 and 6

(C) -6 and 6

(D) -3 and 9

[Ans.D]

Sol. We know that,

$$||z| - |w|| \leq |z - w|$$

$$\Rightarrow -3 < |z| - |w| + 3 < 9.$$

Paragraph for Question Nos. 18 to 20

A circle C of radius 1 is inscribed in an equilateral triangle PQR . The points of contact of C with the sides PQ , QR , RP are D , E , F respectively. The line PQ is given by the equation $\sqrt{3}x + y - 6 = 0$ and the point D is $\left(\frac{3\sqrt{3}}{2}, \frac{3}{2}\right)$. Further, it is given that the origin and the centre of C are on the same side of the line PQ .

18. The equation of circle C is

(A) $(x - 2\sqrt{3})^2 + (y - 1)^2 = 1$

(B) $(x - 2\sqrt{3})^2 + \left(y + \frac{1}{2}\right)^2 = 1$


(C) $(x - \sqrt{3})^2 + (y + 1)^2 = 1$

(D) $(x - \sqrt{3})^2 + (y - 1)^2 = 1$

[Ans.D]

Sol. Let C be the centre of circle and PQ line is $\sqrt{3}x + y - 6 = 0$

By using parametric from you can find centre $(\sqrt{3}, 1)$.



Circle $C : (x - \sqrt{3})^2 + (y - 1)^2 = 1$

19. Points E and F are given by

(A) $\left(\frac{\sqrt{3}}{2}, \frac{3}{2}\right), (\sqrt{3}, 0)$

(B) $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right), (\sqrt{3}, 0)$

(C) $\left(\frac{\sqrt{3}}{2}, \frac{3}{2}\right), \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$

(D) $\left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$

[Ans. A]

Sol. Find the equation of sides making 60° with PQ , those are $y = \sqrt{3}x$ and $y = 0$ and Now you can find the mid points E and F .

20. Equations of the sides QR, RP are

(A) $y = \frac{2}{\sqrt{3}}x + 1, y = -\frac{2}{\sqrt{3}}x - 1$ (B) $y = \frac{1}{\sqrt{3}}x, y = 0$

(C) $y = \frac{\sqrt{3}}{2}x + 1, y = \frac{\sqrt{3}}{2}x - 1$ (D) $y = \sqrt{3}x, y = 0$

[Ans. D]

Sol. Find the equation of sides making 60° with PQ, those are $y = \sqrt{3}x$ and $y = 0$.

Paragraph for Question Nos. 21 to 23

Consider the functions defined implicitly by the equation $y^3 - 3y + x = 0$ on various intervals in the real line. If $x \in (-\infty, -2) \cup (2, \infty)$, the equation implicitly defines a unique real valued differentiable function $y = f(x)$.

If $x \in (-2, 2)$, the equation implicitly defines a unique real valued differentiable function $y = g(x)$ satisfying $g(0) = 0$.

21. If $f(-10\sqrt{2}) = 2\sqrt{2}$, then $f''(-10\sqrt{2}) =$

(A) $\frac{4\sqrt{2}}{7^3 3^2}$ (B) $-\frac{4\sqrt{2}}{7^3 3^2}$ (C) $\frac{4\sqrt{2}}{7^3 3}$ (D) $-\frac{4\sqrt{2}}{7^3 3}$

[Ans. B]

Sol. $3y^2y' - 3y' + 1 = 0$

$$y' = \frac{-1}{3y^2 - 3}$$

$$y'' = \frac{+1}{(3y^2 - 3)^2} 6yy'$$

$$= \frac{6y}{(3y^2 - 3)^2} \cdot \left(\frac{-1}{3y^2 - 3} \right)$$

$$\text{so } f''(-10\sqrt{2}) = \frac{-6.2\sqrt{2}}{(3.8 - 3)^3}$$

$$= \frac{-12\sqrt{2}}{(21)^3} = \frac{-12\sqrt{2}}{7^3 \cdot 3^3} = \frac{-4\sqrt{2}}{7^3 \cdot 3^2}$$

22. The area of the region bounded by the curve $y = f(x)$, the x-axis, and the lines $x = a$ and $x = b$, where $-\infty < a < b < -2$, is-

(A) $\int_a^b \frac{x}{3((f(x))^2 - 1)} dx + b f(b) - a f(a)$


$$(B) - \int_a^b \frac{x}{3((f(x))^2 - 1)} dx + b f(b) - a f(a)$$

$$(C) \int_a^b \frac{x}{3((f(x))^2 - 1)} dx - b f(b) + a f(a)$$

$$(D) - \int_a^b \frac{x}{3((f(x))^2 - 1)} dx - b f(b) + a f(a)$$

[Ans. A]

Sol. The curve is $y^3 - 3y + x = 0$



from the equation of curve : $y' = \frac{-1}{3(y^2 - 1)} \dots(1)$

Required area = Area ABCGDA – Area DCGD

$$= (b - a) f(a) - \left[- \int_{f(b)}^{f(a)} x dy + (f(a) - f(b)) b \right]$$

$$= b f(b) - a f(a) + \int_{f(b)}^{f(a)} x dy$$

Now from equation (1) $x dy = \frac{-x dx}{3(y^2 - 1)}$

Hence the required area is

$$\begin{aligned} \text{Area} &= b f(b) - a f(a) - \int_b^a \frac{x dx}{3(y^2 - 1)} \\ &= b f(b) - a f(a) + \int_a^b \frac{x dx}{3(y^2 - 1)} \\ &= \int_a^b \frac{x dx}{3(f^2(x) - 1)} + b f(b) - a f(a). \end{aligned}$$

23. $\int_{-1}^1 g'(x) dx =$
 (A) $2g(-1)$ (B) 0 (C) $-2g(1)$ (D) $2g(1)$

[Ans. D]

Sol. $\int_{-1}^1 g'(x) dx = (g(x))_{-1}^1 = g(1) - g(-1) = 2g(1)$

as $g(x)$ is odd function in $(-2, 2)$.

Part – II (PHYSICS)

SECTION – I


Straight Objective Type

This section contains 6 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.


24. A spherically symmetric gravitational system of particles has a mass density $\rho = \begin{cases} \rho_0 & \text{for } r \leq R \\ 0 & \text{for } r > R \end{cases}$

where ρ_0 is a constant. A test mass can undergo circular motion under the influence of the gravitational field of particles. Its speed V as a function of distance r ($0 < r < \infty$) from the centre of the system is represented by


(A)




(B)



(C)




(D)



[Ans.C]

Sol.



For $r < R$ $\frac{GMmr}{R^3} = \frac{mv^2}{r} \quad \therefore v^2 \propto r^2$

$\therefore v \propto r$

For $r > R$ $\frac{GMm}{r^2} = \frac{mv^2}{r} \quad \therefore v \propto \frac{1}{\sqrt{r}}$

25. An ideal gas is expanding such that $PT^2 = \text{constant}$. The coefficient of volume expansion of the gas is

- (A) $\frac{1}{T}$ (B) $\frac{2}{T}$ (C) $\frac{3}{T}$ (D) $\frac{4}{T}$

[Ans.C]

Sol. Given $PT^2 = C$, As $PV = nRT$

$\therefore \frac{nRT^3}{V} = C \quad \therefore V = \frac{nRT^3}{C}$


$\frac{dV}{dT} = \frac{3nRT^2}{C} = \frac{3V}{T}$ [Put $T^3 = \frac{CV}{nR}$]

$\left(\frac{dV}{dT}\right)/V = \frac{3}{T}$

26. Two beams of red and violet colours are made to pass separately through a prism (angle of the prism is 60°). In the position of minimum deviation, the angle of refraction will be

- (A) 30° for both the colours (B) greater for the violet colour
(C) greater for the red colour (D) equal but not 30° for both the colours

[Ans. A]



Sol.

For minimum deviation

$$r_1 = r_2 = \frac{A}{2} = 30^\circ$$

Option A is correct

27. Which one of following statements is **Wrong** in the context of X-rays generated from a X-ray tube ?
- (A) Wavelength of characteristic X-rays decreases when the atomic number of the target increases
 - (B) Cut-off wavelength of the continuous X-rays depends on the atomic number of the target
 - (C) Intensity of the characteristic X-rays depends on the electrical power given to the X-ray tube
 - (D) Cut-off wavelength of the continuous X-rays depends on the energy of the electrons in the X-ray tube

[Ans.B]


Sol. (λ_c) cutoff wavelength is given as

$$\lambda_c = \frac{hc}{\text{K.E.}} = \frac{hc}{eV}$$

Hence λ_c does not depend on atomic number of target but depends on potential difference between cathode and anode (target)

28. Figure shows three resistor configurations R1, R2 and R3 connected to 3 V battery. If the power dissipated by the configuration R1, R2 and R3 is P1, P2 and P3, respectively, then

Figure :




- (A) $P_1 > P_2 > P_3$
- (B) $P_1 > P_3 > P_2$
- (C) $P_2 > P_1 > P_3$
- (D) $P_3 > P_2 > P_1$


[Ans.C]

Sol. Resistance of R_1 using wheat stone bridge equivalent circuit

Req. of $R_1 = 1 \Omega$



Equivalent resistance R_2



Similarly R_{eq} of $R_3 = 2\Omega$

$$\text{Power} = \frac{V^2}{R}$$

V is same for all

$$\therefore \text{power} \propto \frac{1}{R}$$

$$\therefore R_2 < R_1 < R_3$$

$$\therefore P_2 > P_1 > P_3$$

29. Students I, II and III perform an experiment for measuring the acceleration due to gravity (g) using a simple pendulum. They use different lengths of the pendulum and/or record time for different number of oscillations. The observations are shown in the table.

Least count for length = 0.1 cm

Least count for time = 0.1 s

Student	Length of the pendulum (cm)	Number of oscillations (n)	Total time for (n) oscillations(s)	Time period(s)
I	64.0	8	128.0	16.0
II	64.0	4	64.0	16.0
III	20.0	4	36.0	9.0

If E_I , E_{II} and E_{III} are the percentage errors in g , i.e. $\left(\frac{\Delta g}{g} \times 100\right)$ for students, I, II, and III respectively,

- (A) $E_I = 0$ (B) E_I is minimum
 (C) $E_I = E_{II}$ (D) E_{II} is maximum

[Ans.B]

Sol.

Student	% error in ℓ (% ℓ)	% error in T (% T)	% error in $g = \% \ell + 2 \times \% T$
I	$\frac{0.1}{64} \times 100 = 0.156$	$\frac{1}{8} \times \frac{0.1}{128} \times 100 = 0.00976$	0.1755
II	$\frac{0.1}{64} \times 100 = 0.156$	$\frac{1}{4} \times \frac{0.1}{64} \times 100 = 0.03906$	0.2341
III	$\frac{0.1}{20} \times 100 = 0.5$	$\frac{1}{4} \times \frac{0.1}{36} \times 100 = 0.0694$	0.6388

SECTION – II

Multiple Correct Answers Type

This section contains 4 multiple correct answer(s) type questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONE OR MORE** is/are correct.

30. Two balls, having linear momenta $\vec{p}_1 = p\hat{i}$ and $\vec{p}_2 = -p\hat{i}$, undergo a collision in free space. There is no external force acting on the balls. Let \vec{p}'_1 and \vec{p}'_2 be their final momenta. The following option(s) is (are) NOT ALLOWED for any non-zero value of p , a_1 , a_2 , b_1 , b_2 , c_1 and c_2 .
- (A) $\vec{p}'_1 = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$; $\vec{p}'_2 = a_2\hat{i} + b_2\hat{j}$ (B) $\vec{p}'_1 = c_1\hat{k}$; $\vec{p}'_2 = c_2\hat{k}$
 (C) $\vec{p}'_1 = a_1\hat{i} + b_1\hat{j} + c_1\hat{k}$; $\vec{p}'_2 = a_2\hat{i} + b_2\hat{j} - c_1\hat{k}$ (D) $\vec{p}'_1 = a_1\hat{i} + b_1\hat{j}$; $\vec{p}'_2 = a_2\hat{i} + b_1\hat{j}$

[Ans.A,D]

Sol. $\vec{P}_1 = P\hat{i}$

$\vec{P}_2 = -P\hat{i}$

Net linear momentum just before collision

$\vec{P}_i = \vec{P}_1 + \vec{P}_2 = P\hat{i} - P\hat{i} = 0$

There is no external force acting on the balls, hence net linear momentum will be conserved.

$P_f = \vec{P}_1' + \vec{P}_2' = 0$

In option (A), $\vec{P}_1' + \vec{P}_2' = (a_1 + a_2)\hat{i} + (b_1 + b_2)\hat{j} + c_1\hat{k} \neq 0$

$\therefore c_1$ is non zero


In option D also

$\vec{P}_1' + \vec{P}_2' = (a_1 + a_2)\hat{i} + 2b_1\hat{j} \neq 0$

$\therefore b_1$ is non zero


- 31.** A particle of mass m and charge q , moving with velocity V enters Region II normal to the boundary as shown in the figure. Region II has a uniform magnetic field B perpendicular to the plane of the paper. The length of the Region II is ℓ . Choose the correct choice(s).

Figure :



- (A) The particle enters Region III only if its velocity $V > \frac{q\ell B}{m}$
- (B) The particle enters Region III only if its velocity $V < \frac{q\ell B}{m}$
- (C) Path length of the particle in Region II is maximum when velocity $V = \frac{q\ell B}{m}$
- (D) Time spent in Region II is same for any velocity V as long as the particle returns to Region I

Sol.




If radius of circle is more than ℓ then particle will be in region III

$$R = \frac{mv}{qB}$$

$$\frac{mv}{qB} > \ell$$


$$v > \frac{q\ell B}{m}$$

Option A is correct option B is wrong



When v is $\frac{qB\ell}{m}$ then particle will take maximum path as shown. Option C is correct.

$$\text{Time period} = \left(\frac{2\pi m}{qB} \right) \frac{1}{2}$$




So this spent time in region II is independent of velocity.

\therefore option (D) is correct.

32. In a Young's double slit experiment, the separation between the two slits is d and the wavelength of the light is λ . The intensity of light falling on slit 1 is four times the intensity of light falling on slit 2. Choose the correct choice(s).
- (A) If $d = \lambda$, the screen will contain only one maximum
- (B) If $\lambda < d < 2\lambda$, at least one more maximum (besides the central maximum) will be observed on the screen
- (C) If the intensity of light falling on slit 1 is reduced so that it becomes equal to that of slit 2, the intensities of the observed dark and bright fringes will increase
- (D) If the intensity of light falling on slit 2 is increased so that it becomes equal to that of slit 1, the intensities of the observed dark and bright fringes will increase


[Ans.A,B]

Sol.



Now when $d = \lambda$

$\Delta x = \lambda$ at ∞ on above side and $\Delta x = -\lambda$ at ∞ on below side. So there are three maxima. One at centre and two are at infinite however screen can't be of infinite size so option A is correct.



$\Rightarrow \lambda < d < 2\lambda \quad \therefore$ say $d = 1.9\lambda$ See above figure

Three maxima are possible as shown the path difference for these maxima are zero, λ and $-\lambda$.

So option B is correct.

Previous intensity of dark fringe = $(\sqrt{4I} - \sqrt{I})^2 = I$


When intensity of both slit become equal then intensity of dark fringe = 0

∴ intensity of dark fringe decreases

∴ option C and D are wrong.

33. Assume that the nuclear binding energy per nucleon (B/A) versus mass number (A) is as shown in the figure. Use this plot to choose the correct choice(s) given below.

Figure :



- (A) Fusion of two nuclei with mass numbers lying in the range of $1 < A < 50$ will release energy
(B) Fusion of two nuclei with mass numbers lying in the range of $51 < A < 100$ will release energy
(C) Fission of a nucleus lying in the mass range of $100 < A < 200$ will release energy when broken into two equal fragments
(D) Fission of a nucleus lying in the mass range of $200 < A < 260$ will release energy when broken into two equal fragments

[Ans. B,D]

- Sol. Energy is released when binding energy per nucleon increases in fusion of two nuclei of mass number from 51 to 100, final nuclei has mass number 102 to 200 where $B.E./A$ is greater similarly in fission of 200 to 260 final mass no becomes 100 to 130.

SECTION – III

Reasoning Type

This section contains 4 reasoning type questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

34. STATEMENT – 1

An astronaut in an orbiting space station above the Earth experiences weightlessness.

and

STATEMENT – 2

An object moving around the Earth under the influence of Earth's gravitational force is in a state of 'free-fall'.

- (A) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is a correct explanation for STATEMENT -1
- (B) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is **NOT** a correct explanation for STATEMENT -1
- (C) STATEMENT-1 is True, STATEMENT-2 is False
- (D) STATEMENT-1 is False, STATEMENT-2 is True

[Ans.A]

Sol.

35. STATEMENT -1

In a Meter Bridge experiment, null point for an unknown resistance is measured. Now, the unknown resistance is put inside an enclosure maintained at a higher temperature. The null point can be obtained at the same point as before by decreasing the value of the standard resistance.


and

STATEMENT-2

Resistance of a metal increases with increase in temperature.

- (A) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is a correct explanation for STATEMENT -1
- (B) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is **NOT** a correct explanation for STATEMENT -1
- (C) STATEMENT-1 is True, STATEMENT-2 is False
- (D) STATEMENT-1 is False, STATEMENT-2 is True

[Ans.D]



Sol.

$$X = R \frac{\ell_1}{\ell_2}$$

As Temperature increase x will increase to get same Null point i.e. same value of $\frac{\ell_1}{\ell_2}$, value of standard

Resistance R should increase.

\therefore Assertion is false, However Reason statement is true

36. STATEMENT – 1

Two cylinders, one hollow (metal) and the other solid (wood) with the same mass and identical dimensions are simultaneously allowed to roll without slipping down an inclined plane from the same height. The hollow cylinder will reach the bottom of the inclined plane first.

and

STATEMENT – 2

By the principle of conservation of energy, the total kinetic energies of both the cylinders are identical when they reach the bottom of the incline.

(A) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is a correct explanation for STATEMENT -1

(B) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is **NOT** a correct explanation for STATEMENT -1

(C) STATEMENT-1 is True, STATEMENT-2 is False

(D) STATEMENT-1 is False, STATEMENT-2 is True

[Ans.D]

Sol. From energy conservation $\frac{1}{2} mv_c^2 + \frac{1}{2} I_c \omega^2 = mgh$

$$\omega = v_c/R$$

$$(I_c)_{\text{solid}} < (I_c)_{\text{hollow}}$$

$$\text{Hence } (v_c)_{\text{solid}} > (v_c)_{\text{hollow}}$$

Hence solid cylinder will reach the bottom first.

37. STATEMENT – 1

The stream of water flowing at high speed from a garden hose pipe tends to spread like a fountain when held vertically up, but tends to narrow down when held vertically down.

and

STATEMENT – 2

In any steady flow of an incompressible fluid, the volume flow rate of the fluid remains constant.

(A) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is a correct explanation for STATEMENT -1

(B) STATEMENT – 1 is True, STATEMENT- 2 is True; STATEMENT -2 is **NOT** a correct explanation

for STATEMENT -1

(C) STATEMENT-1 is True, STATEMENT-2 is False

(D) STATEMENT-1 is False, STATEMENT-2 is True

[Ans.A]

Sol. Volume rate of flow $Q = Av$

When velocity increases cross-sectional area decreases and vice-versa.

SECTION – IV


Linked Comprehension Type

This section contains 3 paragraphs. Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

Paragraph for Question Nos. 38 to 40

A small block of mass M moves on a frictionless surface of an inclined plane, as shown in figure. The angle of the incline suddenly changes from 60° to 30° at point B. The block is initially at rest at A. Assume that collisions between the block and the incline are totally inelastic ($g = 10 \text{ m/s}^2$).

Figure :



38. The speed of the block at point B immediately after it strikes the second incline is -

(A) $\sqrt{60} \text{ m/s}$


(B) $\sqrt{45} \text{ m/s}$

(C) $\sqrt{30} \text{ m/s}$

(D) $\sqrt{15} \text{ m/s}$

[Ans.B]

Sol.



From $\triangle ABD$

$$\tan 60^\circ = \frac{AD}{BD}$$

$$\Rightarrow AD = BD \tan 60^\circ = \sqrt{3} \times \sqrt{3} \text{ m} = 3 \text{ m}$$

speed of block at B just before collision with incline BC is V_B

$$\frac{1}{2} MV_B^2 = Mg(AD) \text{ (Applying conservation of mechanical energy)} \Rightarrow V_B = \sqrt{60} \text{ m/s.}$$

Collision between block and incline is totally inelastic. Just after collision with incline BC component of velocity of block perpendicular to incline BC is zero (because collision is perfectly inelastic)

Component of velocity of block

$$\text{parallel to incline BC is } V'_{B||} = V_B \cos 30^\circ = \sqrt{45} \text{ m/s.}$$

(Component of velocity parallel to surface of contact does not change).

Speed of block just after collision with incline BC is $\sqrt{45} \text{ m/s.}$

39. The speed of the block at point C, immediately before it leaves the second incline is

- (A) $\sqrt{120} \text{ m/s}$ (B) $\sqrt{105} \text{ m/s}$ (C) $\sqrt{90} \text{ m/s}$ (D) $\sqrt{75} \text{ m/s}$

[Ans.B]

Sol. $BD = 3\sqrt{3} \tan 30^\circ = 3 \text{ m}$

From conservation of mechanical energy

Total mechanical energy at point B = total mechanical energy at point C


$$\frac{1}{2} M(\sqrt{45})^2 + Mg(3) = \frac{1}{2} MV_C^2$$

$$V_C = \sqrt{105} \text{ m/s}$$

40. If collision between the block and the incline is completely elastic, then the vertical (upward) component of the velocity of the block at point B, immediately after it strikes the second incline is
- (A) $\sqrt{30}$ m/s (B) $\sqrt{15}$ m/s (C) 0 (D) $-\sqrt{15}$ m/s

[Ans.C]

Sol. Collision is elastic.



Just after collision with incline BC,

$$\text{Component of velocity of block along BC is } V''_{B\parallel} = V_B \cos 30^\circ = \sqrt{45} \text{ m/s}$$

$$\text{Component of velocity perpendicular to BC is } V''_{B\perp r} = V_B \sin 30^\circ = \sqrt{15} \text{ m/s along BF}$$

V''_B = velocity of block just after collision

$$\tan \theta = \frac{V''_{B\perp r}}{V''_{B\parallel}} = \frac{\sqrt{15}}{\sqrt{45}} = \frac{1}{\sqrt{3}}$$

$$\theta = 30^\circ$$


\vec{V}''_B is in horizontal direction

Hence vertical component of \vec{V}''_B is zero.

Paragraph for Question Nos. 41 to 43

A small spherical monoatomic ideal gas bubble ($\gamma = 5/3$) is trapped inside a liquid of density ρ_l (see figure). Assume that the bubble does not exchange any heat with the liquid. The bubble contains n moles of gas. The temperature of the gas when the bubble is at the bottom is T_0 , the height of the liquid is H and the atmospheric pressure is P_0 (Neglect surface tension)


Figure :



41. As the bubble moves upwards, besides the buoyancy force the following forces are acting on it
- (A) Only the force of gravity
- (B) The force due to gravity and the force due to the pressure of the liquid
- (C) The force due to gravity, the force due to the pressure of the liquid and the force due to viscosity of the liquid
- (D) The force due to gravity and the force due to viscosity of the liquid

[Ans.D]

Sol. Free Body Diagram of gas bubble



Buoyancy force is due to pressure difference.

42. When the gas bubble is at a height y from the bottom, its temperature is

(A) $T_0 \left(\frac{P_0 + \rho_\ell gH}{P_0 + \rho_\ell gy} \right)^{2/5}$

(B) $T_0 \left(\frac{P_0 + \rho_\ell g(H-y)}{P_0 + \rho_\ell gH} \right)^{2/5}$

(C) $T_0 \left(\frac{P_0 + \rho_\ell gH}{P_0 + \rho_\ell gy} \right)^{3/5}$


(D) $T_0 \left(\frac{P_0 + \rho_\ell g(H-y)}{P_0 + \rho_\ell gH} \right)^{3/5}$

[Ans.B]

Sol. As the number of moles remains conserved

$$n_i = n_f$$

$$\frac{P_i V_i}{T_i} = \frac{P_f V_f}{T_f}$$



$$T_f \frac{P_f V_f}{P_i V_i} = T_i \quad \dots (1)$$

Since bubble does not exchange heat hence process is adiabatic in nature

$$P_i V_i^\gamma = P_f V_f^\gamma$$

$$V_f/V_i = (P_i/P_f)^{1/\gamma} \quad \dots (2)$$

Putting (2) in (1)

$$T_f = \left(\frac{P_f}{P_i} \right) \left(\frac{P_i}{P_f} \right)^{1/\gamma} \cdot T_i$$

$$T_f = \left(\frac{P_f}{P_i} \right)^{1-1/\gamma} \cdot T_i$$

$$P_i = P_0 + \rho_L g H$$

$$P_f = P_0 + \rho_L g (H - y)$$

$$T_f = \left[\frac{P_0 + \rho_L g (H - y)}{P_0 + \rho_L g H} \right]^{2/5} T_0$$

43. The buoyancy force acting on the gas bubble is (Assume R is the universal gas constant)

$$(A) \rho_L n R g T_0 \frac{(P_0 + \rho_L g H)^{2/5}}{(P_0 + \rho_L g y)^{7/5}}$$

$$(B) \frac{\rho_L n R g T_0}{(P_0 + \rho_L g H)^{2/5} [P_0 + \rho_L g (H - y)]^{3/5}}$$

$$(C) \rho_L n R g T_0 \frac{(P_0 + \rho_L g H)^{3/5}}{(P_0 + \rho_L g y)^{8/5}}$$

$$(D) \frac{\rho_L n R g T_0}{(P_0 + \rho_L g H)^{3/5} [P_0 + \rho_L g (H - y)]^{2/5}}$$

[Ans.B]

Sol. Buoyant force

$$F_B = \rho_L V_f g$$

$$F_B = \rho_L \left(\frac{n R T_f}{P_f} \right) g$$

$$= \frac{\rho_L n R}{P_f} \left(\frac{P_f}{P_i} \right) \left(\frac{P_i}{P_f} \right)^{1/\gamma} T_i$$

$$= \frac{\rho_L n R}{P_i^{1-1/\gamma}} \cdot \frac{1}{P_f^{1/\gamma}} \cdot T_i$$

$$F_B = \frac{\rho_L n R}{P_i^{2/5} P_f^{3/5}} \cdot T_i$$

$$F_B = \frac{\rho_L n R}{(P_0 + \rho_L g H)^{2/5} [P_0 + \rho_L g (H - y)]^{3/5}} T_0$$


Paragraph for Question Nos. 44 to 46

In a mixture of H – He⁺ gas (He⁺ is singly ionized He atom), H atoms and He⁺ ions are excited to their respective first excited states. Subsequently, H atoms transfer their total excitation energy to He⁺ ions (by collisions). Assume that the Bohr model of atom is exactly valid.

44. The quantum number n of the state finally populated in He^+ ions is
 (A) 2 (B) 3 (C) 4 (D) 5

[Ans.C]


Sol. Energy level diagram



H-atom

(First excited state)

When hydrogen atom transfers its total excitation energy to He^+ ion then He^+ ion are excited to $n = 4$ quantum state as seen from energy level diagram



He^+ ion

First excited state

Hence quantum number of state finally populated in He^+ ions is 4.

45. The wavelength of light emitted in the visible region by He^+ ions after collisions with H atoms is
 (A) 6.5×10^{-7} m (B) 5.6×10^{-7} m
 (C) 4.8×10^{-7} m (D) 4.0×10^{-7} m

[Ans.C]

Sol. The wavelength of light emitted in the visible region by He^+ ions in final excited state is

$$\lambda = \frac{hc}{\Delta E_{4 \rightarrow 3}} \quad \Delta E_{4 \rightarrow 3} = (6.04 - 3.4) \text{ eV}$$

$$= 2.64 \text{ eV} \quad \lambda = \frac{hc}{2.64 \text{ eV}}$$

$$\lambda = \frac{12.4 \times 10^{-7}}{2.64} \quad l = 4.8 \times 10^{-7} \text{ m}$$

46. The ratio of the kinetic energy of the $n = 2$ electron for the H atom to that of He^+ ion is

- (A) $\frac{1}{4}$ (B) $\frac{1}{2}$ (C) 1 (D) 2

[Ans.A]

Sol. $\text{K.E.} \propto Z^2/n^2$

$$\frac{\text{K.E.}_H}{\text{K.E.}_{\text{He}^+}} = \frac{Z_H^2}{Z_{\text{He}^+}^2} = \frac{(1)}{(2)^2} = \frac{1}{4}$$

Part – III (CHEMISTRY)

SECTION – I

Straight Objective Type

This section contains 6 multiple choice questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

47. 2.5 mL of $\frac{2}{5}$ M weak monoacidic base ($K_b = 1 \times 10^{-12}$ at 25°C) is titrated with $\frac{2}{15}$ M HCl in water at 25°C . The concentration of H^+ at equivalence point is ($K_w = 1 \times 10^{-14}$ at 25°C)
- (A) 3.7×10^{-13} M (B) 3.2×10^{-7} M
 (C) 3.2×10^{-2} M (D) 2.7×10^{-2} M

[Ans. D]

Sol. Moles of BOH taken = $2.5 \times \frac{2}{5} \times 10^{-3}$, = 10^{-3} moles

Moles of HCl required = 10^{-3} moles.

$$V \times \frac{2}{15} \times 10^{-3} = 10^{-3}$$

\therefore Vol. of HCl required $V = 15/2$ mL

Total vol. = $7.5 + 2.5 = 10$ ml

moles of BCl obtained = 10^{-3} moles.

$$[\text{BCl}] = \frac{10^{-3}}{10} \times 10^3 = 0.1 \text{ (M)}$$

$$\frac{Ch^2}{(1-h)} = \frac{K_w}{K_b} = \frac{10^{-14}}{10^{-12}} = 10^{-2} \quad \text{or} \quad h \approx 0.27$$

$$0.1 \times \frac{h^2}{(1-h)} = 10^{-2}$$

$$[H^+] = Ch = 2.7 \times 10^{-2}$$

48. Under the same reaction conditions, initial concentration of $1.386 \text{ mol dm}^{-3}$ of a substance becomes half in 40 seconds and 20 seconds through first order and zero order kinetics, respectively. Ratio $\left(\frac{k_1}{k_0}\right)$ of the rate constants for first order (k_1) and zero order (k_0) of the reactions is -
- (A) $0.5 \text{ mol}^{-1} \text{ dm}^3$ (B) 1.0 mol dm^{-3}
 (C) 1.5 mol dm^{-3} (D) $2.0 \text{ mol}^{-1} \text{ dm}^3$ [Ans. A]

Sol.

$$K_1 = \frac{0.693}{t_{1/2}} = \frac{0.693}{40} \text{ s}^{-1}$$

$$-\frac{d[A]}{dt} = K_0$$

$$-d[A] = K_0 dt$$

$$[A]_0 - [A] = K_0 t$$

$$[A]_0 - \frac{[A]_0}{2} = K_0 t_{1/2}$$

$$K_0 = \frac{[A]_0}{2t_{1/2}} = \frac{[A]_0}{2 \times 20} = \frac{1.386}{2 \times 20} \text{ mole lit}^{-1} \text{ s}^{-1}$$

$$\frac{K_1}{K_0} = \frac{\frac{0.693}{40}}{\frac{1.386}{2 \times 20}} = \frac{0.693}{1.386} \times \frac{40}{2} = \frac{1}{2} \frac{\text{s}^{-1}}{\text{mol dm}^{-3} \text{ s}^{-1}} = 0.5 \text{ dm}^3 \text{ mol}^{-1}$$

49. Native silver metal forms a water soluble complex with a dilute aqueous solution of NaCN in the presence of -
- (A) nitrogen (B) oxygen
 (C) carbon dioxide (D) argon [Ans. B]

Sol.




50. Aqueous solution of $\text{Na}_2\text{S}_2\text{O}_3$ on reaction with Cl_2 gives -

- (A) $\text{Na}_2\text{S}_4\text{O}_6$ (B) NaHSO_4
 (C) NaCl (D) NaOH

[Ans. C]

Sol. $4 \text{Na}_2\text{S}_2\text{O}_3 + \text{Cl}_2 \rightarrow 3 \text{Na}_2\text{SO}_4 + 2 \text{NaCl} + 5\text{S}\downarrow$


51. The major product of the following reaction is -



- (A)
- (B)
- (C)
- (D)

[Ans. A]

Sol.



In SN^2 mech., Inversion takes place at alkyl position.

52. Hyperconjugation involves overlap of the following orbitals -

- (A) $\sigma - \sigma$ (B) $\sigma - p$ (C) $p - p$ (D) $\pi - \pi$


[Ans. B]

SECTION – II

Multiple Correct Answers Type

This section contains 4 multiple correct answer(s) type questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONE OR MORE** is/are correct.


53. The correct statement(s) about the compound given below is (are)



- (A) The compound is optically active
(B) The compound possesses centre of symmetry
(C) The compound possesses plane of symmetry
(D) The compound possesses axis of symmetry

[Ans. A]

- Sol. Fisher projection of given structure will be



This is threo form & optically active without any element of symmetry.

54. A solution of colourless salt H on boiling with excess NaOH produces a non-flammable gas. The gas evolution ceases after sometime. Upon addition of Zn dust to the same solution, the gas evolution restarts. The colourless salt (s) H is (are) -
- (A) NH_4NO_3 (B) NH_4NO_2 (C) NH_4Cl (D) $(\text{NH}_4)_2\text{SO}_4$

[Ans. A, B]

- Sol. $\text{NH}_4\text{NO}_3 + \text{NaOH} \rightarrow \text{NaNO}_3 + \text{NH}_3 \uparrow + \text{H}_2\text{O}$
 $\text{Zn} + 2 \text{NaOH} \rightarrow \text{Na}_2\text{ZnO}_2 + 2\text{H}$
 $\text{NaNO}_3 + 8\text{H} \rightarrow \text{NaOH} + \text{NH}_3 \uparrow + 2\text{H}_2\text{O}$
 $\text{NH}_4\text{NO}_2 + \text{NaOH} \rightarrow \text{NaNO}_2 + \text{NH}_3 \uparrow + \text{H}_2\text{O}$
 $\text{NaNO}_2 + 6\text{H} \rightarrow \text{NaOH} + \text{NH}_3 \uparrow + \text{H}_2\text{O}$

55. A gas described by van der Waals equation -
- (A) behaves similar to an ideal gas in the limit of large molar volumes
 (B) behaves similar to an ideal gas in the limit of large pressures
 (C) is characterised by van der Waals coefficients that are dependent on the identity of the gas but are independent of the temperature
 (D) has the pressure that is lower than the pressure exerted by the same gas behaving ideally

[Ans. A, D]


Sol. Due to inter molecular force real gas has lesser pressure than ideal gas.

$$\left(P + \frac{a}{V_m^2} \right) (V_m - b) = RT$$


When $V_m \rightarrow \infty$

$$PV_m \approx RT$$


56. The correct statement(s) concerning the structures E, F and G is (are)



(E)



(F)



(G)

- (A) E, F and G are resonance structures
 (B) E, F and E, G are tautomers
 (C) F and G are geometrical isomers
 (D) F and G are diastereomers

[Ans. B, C, D]

SECTION – III

Reasoning Type

This section contains 4 reasoning type questions. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

57. **STATEMENT-1** : Pb^{4+} compounds are stronger oxidizing agents than Sn^{4+} compounds.
and
STATEMENT-2 : The higher oxidation states for the group 14 elements are more stable for the heavier members of the group due to inert pair effect.
- (A) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is a correct explanation for STATEMENT-1.
 (B) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is **NOT** a correct explanation for STATEMENT-1.

(C) STATEMENT-1 is True, STATEMENT-2 is False.

(D) STATEMENT-1 is False, STATEMENT-2 is True.

[Ans. C]

58. **STATEMENT-1** : Bromobenzene upon reaction with Br_2/Fe gives 1,4-dibromobenzene as the major product.

and

STATEMENT-2 : In bromobenzene, the inductive effect of the bromo group is more dominant than the mesomeric effect in directing the incoming electrophile.

(A) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is a correct explanation for STATEMENT-1.

(B) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is **NOT** a correct explanation for STATEMENT-1.

(C) STATEMENT-1 is True, STATEMENT-2 is False.

(D) STATEMENT-1 is False, STATEMENT-2 is True.

[Ans. C]

Sol. Due to halogen substituent reactivity depends more on I effect & orientation of attack depends more on +M effect.

59. **STATEMENT-1** : The plot of atomic number (y-axis) versus number of neutrons (x-axis) for stable nuclei shows a curvature towards x-axis from the line of 45° slope as the atomic number is increased.

and

STATEMENT-2 : Proton-proton electrostatic repulsions begin to overcome attractive forces involving protons and neutrons in heavier nuclides.

(A) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is a correct explanation for STATEMENT-1.


(B) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is **NOT** a correct explanation for STATEMENT-1.

(C) STATEMENT-1 is True, STATEMENT-2 is False.

(D) STATEMENT-1 is False, STATEMENT-2 is True.

[Ans. A]

Sol.



With increasing Z, increase in n outweighs increase in Z

60. **STATEMENT-1** : For every chemical reaction at equilibrium, standard Gibbs energy of reaction is zero.
and

STATEMENT-2 : At constant temperature and pressure, chemical reactions are spontaneous in the direction of decreasing Gibbs energy.

- (A) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is a correct explanation for STATEMENT-1.
 (B) STATEMENT-1 is True, STATEMENT-2 is True; STATEMENT-2 is **NOT** a correct explanation for STATEMENT-1.
 (C) STATEMENT-1 is True, STATEMENT-2 is False.
 (D) STATEMENT-1 is False, STATEMENT-2 is True.

[Ans. D]

Sol. At equilibrium $\Delta_r G = 0$ & $\Delta_r G^\circ$ may or may not be zero.

SECTION – IV

Linked Comprehension Type

This section contains 3 paragraphs. Based upon each paragraph, 3 multiple choice questions have to be answered. Each question has 4 choices (A), (B), (C) and (D), out of which **ONLY ONE** is correct.

Paragraph for Question Nos. 61 to 63

Properties such as boiling point, freezing point and vapour pressure of a pure solvent change when solute molecules are added to get homogeneous solution. These are called colligative properties. Applications of colligative properties are very useful in day-to-day life. One of its examples is the use of ethylene glycol and water mixture as anti-freezing liquid in the radiator of automobiles.

A solution **M** is prepared by mixing ethanol and water. The mole fraction of ethanol in the mixture is 0.9.

Given : Freezing point depression constant of water (K_f^{water}) = $1.86 \text{ K kg mol}^{-1}$

Freezing point depression constant of ethanol ($K_f^{\text{ethanol}} = 2.0 \text{ K kg mol}^{-1}$)

Boiling point elevation constant of water ($K_b^{\text{water}} = 0.52 \text{ K kg mol}^{-1}$)

Boiling point elevation constant of ethanol ($K_b^{\text{ethanol}} = 1.2 \text{ K kg mol}^{-1}$)

Standard freezing point of water = 273 K

Standard freezing point of ethanol = 373 K

Standard boiling point of ethanol = 351.5 K

Vapour pressure of pure water = 32.8 mm Hg

Vapour pressure of pure ethanol = 40 mm Hg

Molecular weight of water = 18 g mol⁻¹

Molecular weight of ethanol = 46 g mol⁻¹

In answering the following questions, consider the solutions to be ideal dilute solutions and solutes to be non-volatile and non-dissociative.

61. The freezing point of the solution **M** is -

(A) 268.7 K

(B) 268.5 K

(C) 234.2 K

(D) 150.9 K

[Ans. D]

Sol. $x_{\text{C}_2\text{H}_5\text{OH}} = 0.9$

$x_{\text{H}_2\text{O}} = 0.1$

In $0.9 \times 46 \text{ gm C}_2\text{H}_5\text{OH}$, H_2O present = 0.1 mole

in 1000 gm $\text{C}_2\text{H}_5\text{OH}$, H_2O present = $\frac{0.1 \times 1000}{0.9 \times 46} = 2.415 \text{ (m)}$

$\Delta T_f = K_f \times m = 2 \times 2.415 = 4.83$

\therefore Freezing point of solⁿ. = $155.7 - 4.83 = 150.87 \text{ K}$

62. The vapour pressure of the solution **M** is -

(A) 39.3 mm Hg

(B) 36.0 mm Hg

(C) 29.5 mm Hg

(D) 28.8 mm Hg

[Ans. A]

Sol. $P_{\text{total}} = x_{\text{C}_2\text{H}_5\text{OH}} \cdot P_{\text{C}_2\text{H}_5\text{OH}}^{\circ} + x_{\text{H}_2\text{O}} \cdot P_{\text{H}_2\text{O}}^{\circ}$

$P_{\text{total}} = 0.9 \times 40 + 0.1 \times 32.8$ $P_{\text{total}} = 39.28 \text{ torr}$

63. Water is added to the solution **M** such that the mole fraction of water in the solution becomes 0.9. The boiling point of this solution is -

(A) 380.4 K

(B) 376.2 K

(C) 375.5 K

(D) 354.7 K

[Ans. B]

Sol. In this case water is a solvent and $\text{C}_2\text{H}_5\text{OH}$ is a solute

$x_{\text{C}_2\text{H}_5\text{OH}} = 0.1$

$x_{\text{H}_2\text{O}} = 0.9$

In 0.9×18 gm H_2O , C_2H_5OH present = 0.1 mole

In 1000 gm H_2O , C_2H_5OH present = $\frac{0.1 \times 1000}{0.9 \times 18} = 6.17$ (molality)


$\Delta T_b = K_b \times m = 0.52 \times 6.17 = 3.2084$

\therefore Boiling point of the solution




= $373 + 3.2084 = 376.2$ K

Paragraph for Questions Nos. 64 to 66

In the following reaction sequence, products I, J and L are formed. K represents a reagent.







64. The structure of the product I is -






[Ans. D]

65. The structures of compounds J and K, respectively, are -




[Ans. A]

66. The structure of product L is -




[Ans. C]

Sol. 64 to 66



[I]



Paragraph for Question Nos. 67 to 69

There are some deposits of nitrates and phosphates in earth's crust. Nitrates are more soluble in water. Nitrates are difficult to reduce under the laboratory conditions but microbes do it easily. Ammonia forms large number of complexes with transition metal ions. Hybridization easily explains the ease of sigma donation capability of NH_3 and PH_3 . Phosphine is a flammable gas and is prepared from white phosphorous.

67. Among the following, the correct statement is -
- (A) Phosphates have no biological significance in humans
 - (B) Between nitrates and phosphates, phosphates are less abundant in earth's crust
 - (C) Between nitrates and phosphates, nitrates are less abundant in earth's crust
 - (D) Oxidation of nitrates is possible in soil
- [Ans. C]**
68. Among the following, the correct statement is -
- (A) Between NH_3 and PH_3 , NH_3 is a better electron donor because the lone pair of electrons occupies spherical 's' orbital and is less directional
 - (B) Between NH_3 and PH_3 , PH_3 is a better electron donor because the lone pair of electrons occupies sp^3 orbital and is more directional
 - (C) Between NH_3 and PH_3 , NH_3 is a better electron donor because the lone pair of electrons occupies sp^3 orbital and is more directional

(D) Between NH_3 and PH_3 , PH_3 is a better electron donor because the lone pair of electrons occupies spherical 's' orbital and is less directional

[Ans. C]

Sol. In PH_3 p-character is mostly used for bonding while lone pair is in predominantly s-character orbital which is diffused & less reactive.

69. White phosphorus on reaction with NaOH gives PH_3 as one of the products. This is a -

(A) dimerization reaction

(B) disproportionation reaction

(C) condensation reaction

(D) precipitation reaction

[Ans. B]

